


New to Travel Course – Day 1

Friday 6th September 2019
Written and taught by
Jane Chiodini MSc RGM RM FFTM RCPS(Glasg)
Queen's Nurse



Jane Chiodini
Travel Health Specialist Nurse

1

About you




Where from?
Your job?
Your travel health experience?

Jane Chiodini
Travel Health Specialist Nurse

2

What do you hope to achieve from this course?




Write two goals down on the form provided

Jane Chiodini
Travel Health Specialist Nurse

3

What are my goals



1. **Today** - I want you to understand the basic principles of a risk assessment, the diseases and vaccines and the resources to help put safe service together
2. **By the end of the two days** I'd like you to go away feeling more 'in control' for travel health, hopefully enthused about the subject and to potentially enjoy it in the future!

Competence comes with time and experience!

Jane Chiodini
Travel Health Specialist Nurse

4

www.janechiodini.co.uk/education/new-to-travel/sept2019



Jane Chiodini
Travel Health Specialist Nurse

5

Agenda

- Introduction to travel medicine
- Travel risk assessment
- Travel vaccines and related issues
- Travel medicine operational issues
- Recap on resources

Jane Chiodini
Travel Health Specialist Nurse

6

Recap of the subject onlinelearning

<http://www.janechiodini.co.uk/education/online-learning/>

Jane Chiodini
Travel Health Specialist Nurse

7

An introduction to travel medicine – the key issues

- The travelling public needs to be well informed not only about their destinations and all of the cultural richness, but also **aware of the potential risks during their journey**
- Equally there needs to be a nucleus of GPs, practice nurses and other trained health professionals who are **knowledgeable about the risks on a country by country basis and who are confident about advising their patients** about each of the measures necessary to keep them healthy while travelling

Field VE, Ford L, Hill DR, eds. Health Information for Overseas Travel, National Travel Health Network and Centre, London, UK, 2010.

Jane Chiodini
Travel Health Specialist Nurse

8

Who provides travel health advice?

- In other countries its traditionally the doctor who sees the traveller and performs the risk assessment, passing them on to the nurse to administer the vaccines and give some advice
- More recently pharmacists have become involved in travel medicine, especially in Canada and the UK
- In the UK, nurses have been undertaking all aspects of travel health since the early 1990s, from risk assessment to administration of vaccines and providing risk management advice. In some cases, nurses who have obtained a non medical prescribing qualification are not only prescribing but in some circumstances setting up and owning their own travel clinics

Jane Chiodini
Travel Health Specialist Nurse

9

Why is the practice of travel medicine different in the UK?

- National Health Service provides some travel vaccines free of charge – service provided in the majority of primary care settings as GPs are financially rewarded for the service
- Pressure on GPs with their workload so historically, they passed travel health on to the nurses, but now pharmacists are getting very involved as well, with private clinics are growing dramatically
- Some surgeries are ceasing the provision of a travel service – however, they are NOT allowed to do this unless they surrender the provision of the global sum which they receive for immunisation services

Jane Chiodini
Travel Health Specialist Nurse

10

www.janechiodini.co.uk/news/faqs/faq-2/

Jane Chiodini
Travel Health Specialist Nurse

Jane Chiodini
Travel Health Specialist Nurse

11

Charging for travel vaccines in an NHS setting – covered in day 2

- Vaccines that must always be given as part of NHS provision** (hepatitis A all doses, combination A+B all doses, typhoid, combination typhoid and hep A, polio and cholera)
- Vaccines that cannot be given as an NHS service** (yellow fever, Japanese encephalitis, tick borne encephalitis and rabies for travel and more recently ACWY for travel – but see FAQ page)
- Vaccines that can be given as NHS or private service** (hepatitis B)


Note: Cholera and oral typhoid vaccines are now only NHS vaccines in an NHS setting

see <http://www.janechiodini.co.uk/news/faqs/faq-no-2/>

Jane Chiodini
Travel Health Specialist Nurse

12

General Practice in the UK



Bedfordshire

UK GP Statistics 2017¹


9,800 practices

- 7,613 in England
- 958 in Scotland
- 454 in Wales
- 349 in Northern Ireland

51,000+ GPs

New data from NHS²

Digital reported 22,976 practice nurses working in primary care as of March 2018 with a full-time equivalent of 15,889






1. BMA Press Briefing – General Practice in the UK July 2017 (accessed January 2016)
<https://www.bma.org.uk/media/files/pdfs/news%20views%20analysis/press%20briefings/general-practice.pdf>
<https://www.nursingpractice.com/full-time-practice-nurse-numbers-increase-more-2>

Jane Chiodini
 Travel and Clinical Medicine

13

Governance for health care professionals – working within ‘our codes’






http://www.gmc-uk.org/Good_medical_practice_English_1215.pdf_51527435.pdf
<https://www.pharmacyregulation.org/page>
http://www.nmc-uk.org/Documents/Standards/The_code-M-20100006.pdf

Jane Chiodini
 Travel and Clinical Medicine

14

A key document for those undertaking travel medicine



Includes

- History of travel medicine
- Details about the provision of a travel service
- Risk assessment
- Competencies
- Forms
- Resources

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN.

Jane Chiodini
 Travel and Clinical Medicine

15

New inclusions - Page 9

A statement is included for those who run Yellow Fever Vaccination Centres in the UK acknowledging that whilst YF training is not mandatory for all individuals administering the vaccine, both NaTHNaC and Health Protection Scotland (HPS) recommend:

all those responsible for administering YF vaccine complete the training for their own accountability and good practice

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN.

Jane Chiodini
 Travel and Clinical Medicine

16

Three levels of nurses


Competent nurse	Experienced / proficient nurse	Senior practitioner / expert nurse
See slides to follow outlining expectations	Fulfils points of competent nurse as well	Fulfils points of competent and experienced nurse as well

While there is a strong focus on the work of a registered nurse, the field of travel medicine is truly multidisciplinary and much of the information provided in this publication is equally applicable to other registered health care professionals including **doctors** and **pharmacists** who provide travel health

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN.

Jane Chiodini
 Travel and Clinical Medicine

17



For doctors, pharmacists and nurses specialising in travel medicine

Go to
<http://www.janechiodini.co.uk/about/publications/>

<http://download.journals.elsevierhealth.com/pdfs/journals/1477-8939/PIIS1477893912000671.pdf>

Jane Chiodini
 Travel and Clinical Medicine

18

Core competence for the *Competent Nurse* (or practitioner) in a travel health consultation (pages 21/23)


- Demonstrates good geographical knowledge
- Able to perform risk assessment effectively and understands how to interpret potential risk within a trip
- Knows where to 'go' for recommendations for travel advice, immunisations, malaria chemoprophylaxis
- Recognises limit of knowledge and knows when to refer appropriately
- Has good knowledge of common travel related illnesses e.g. TD, hepatitis, typhoid, malaria

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse

19

Search on 'travel health' at www.nmc.org.uk/

i) Documented in the patient record that the patient was going to 'Lybia' (sic.) (a reference to Libya, a low risk travel destination for contracting malaria) when in fact the patient was travelling to Liberia (a high risk travel destination for contracting malaria).



<https://goo.gl/SqtoQF>


Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse

20

Core competence continued

Able to provide individual advice to the traveller

- ✓ Accident prevention
- ✓ Safe food, water and personal hygiene
- ✓ Prevention of blood-borne infections and sexually transmitted diseases
- ✓ General insect bite prevention
- ✓ Prevention of animal bites, particularly rabies including wound management
- ✓ Prevention of sun and heat complications
- ✓ Personal safety and security
- ✓ Malaria awareness, ABCD advice



Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse

21

Core competence continued

- Communicates information effectively
- Prioritises in a situation when traveller is on a limited budget
- Assesses anxieties and acts appropriately
- Demonstrates an excellent vaccine administration technique
- Completes patient and administrative records after vaccination

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse

22

Education and Training – page 23

- Demonstrates evidence of learning to apply skills and knowledge in the field of travel medicine. For example, minimum of 15 hours of relevant learning plus mentorship in clinical skills before undertaking a travel consultation alone
- Ensures travel health knowledge is always up to date
- Attends an annual travel health update study session/conference at a local, national or international event

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse

23

Agenda

- ✓ Introduction to travel medicine
 - Travel risk assessment
 - Travel vaccines and related issues
 - Travel medicine operational issues
 - Recap on resources

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse

24

Risk Assessment & Management in Travel Health



Jane Chiodini
Travel Health Consultant

25

Aims and Objectives of this session

- To understand what risk assessment is
- To appreciate the elements of the risk assessment process
- To have a good understanding of the required knowledge and resources needed to perform a risk assessment
- To be able to apply these skills at the end of the course

Jane Chiodini
Travel Health Consultant


26

Risk assessment

No travel health consultation should take place without conducting a travel risk assessment and documenting the information.

The assessment forms the basis of all subsequent decisions, advice given, vaccines administered and the malaria prophylaxis advice that is offered.

This takes time to perform correctly, and for best practice practitioners should leave sufficient time.



Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN.

Jane Chiodini
Travel Health Consultant

27

Time is the major constraint

From the RCN Guidance – page 18

The main consideration is to allocate sufficient time to perform the risk assessment. It would be **unsafe to only allow 10 - 15 minutes** for a new travel appointment.

A 20-minute consultation appointment per person should be allowed to exercise best practice. Travellers with more complex needs such as backpackers or individuals requiring malaria prevention advice relevant to their destination - **may need even longer** consultation time.

The Nursing and Midwifery Council 'Code' is about being professional, about being accountable and about being able to justify your decisions; employers need to respect the complexity of a travel consultation and appreciate that sufficient time must be allowed for nurses to abide by the Code.

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN.

Jane Chiodini
Travel Health Consultant

28


What is risk?



Jane Chiodini
Travel Health Consultant

29

Would you enjoy this?



Jane Chiodini
Travel Health Consultant

30

The best way to manage/conduct a risk assessment – one option

Travel risk assessment form completed prior to appointment by traveller

Travel risk assessment form reviewed by travel health adviser

Management of the travel risks discussed with the traveller by the travel health adviser and conclusions reached

Jane Chiodini
Travel health adviser

31

Example of risk assessment form for information recording

Available to download from my 'Tools' page – item no. 1 <http://www.janechiodini.co.uk/tools/>

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN.

Jane Chiodini
Travel health adviser

32

Information to be gathered

Traveller information	Traveller's itinerary
<ul style="list-style-type: none"> Age and sex Medical history, past and present Current health status Medication Allergies to drugs and food Previous experience travelling Current interest and knowledge of health risks Previous vaccine history Any special needs 	<ul style="list-style-type: none"> Destinations (s) Date of departure Duration of stay Mode of transport Purpose of trip and planned activities Quality of accommodation Financial budget Healthcare standards at destination Relevant comprehensive insurance provision

Jane Chiodini
Travel health adviser

33

Risk assessment exercise

- Beckham is 10 years old and is travelling to Angola in the summer holidays to stay with his grandparents for 8 weeks
- What are the issues and risks when assessing this traveller?

What if Beckham had been a girl, is there anything else you might consider?

Jane Chiodini
Travel health adviser

34

Risk management

Having performed a risk assessment the risks identified are managed by individualised advice

<ul style="list-style-type: none"> Medical preparation Journey risks Safety risks Environmental risks Food and water borne risks 	<ul style="list-style-type: none"> Vector borne risks Air borne risks Sexual health and blood borne viral risks Skin health Psychological health
---	---

<https://travelhealthpro.org.uk/>

Jane Chiodini
Travel health adviser

35

What does performing a risk assessment achieve?

It enables you to give:

<p>Appropriate travel health risk advice</p>	<p>Appropriate travel vaccines for travel plans</p>	<p>Appropriate malaria prevention advice</p>
--	---	--

To perform and provide evidence of best practice

Jane Chiodini
Travel health adviser

36

Implications of the questions we ask

Many sources to increase your knowledge and understanding of pre-travel risk assessment in more detail – including on national databases and international resources



Jane Chiodini
Travel Health Consultant

44

The following slides provide some examples
but please refer to the resources on previous slide for more information.

Jane Chiodini
Travel Health Consultant

45

Age and Sex



Jane Chiodini
Travel Health Consultant

46

Babies and small children


- Increased risk of other hazards e.g. accidents, encounters with animals – need for rabies post exposure
- Small, mobile, inquisitive toddlers, limited hygiene awareness
- Risk of illness more severe – e.g travellers' diarrhoea, malaria – requiring medical treatment abroad
- Restrictions on some choices of vaccines and malaria chemoprophylaxis

Jane Chiodini
Travel Health Consultant

47

Older travellers

- Immune systems reduced – infection risk increased
- Senses reduced
- PMH more common
- Immunisation status
- Specific problems e.g. yellow fever vaccine




Jane Chiodini
Travel Health Consultant

48

Female travellers

- Security risk
- Travelling during pregnancy / breast feeding
- Managing contraception
- Coping with menstruation



Jane Chiodini
Travel Health Consultant

49

www.mooncup.co.uk

mooncup® an innovative alternative to tampons

home
what is it?
why is it better?
is it easy to use?
which size?
price
get convinced?
how to use
faq
history
testimonials
order online
wh stockists
contact us

newsletter
tell a friend
wholesale info
sell the mooncup
resources

© Mooncup Ltd 2005

"3 days after using my mooncup for the first time and I want to tell the world what they are missing out on! I keep forgetting to wear one this month. I was dubious at first but now I love it and am recommending it to all my friends!"

Angie

- What is it?
- What are women saying about the Mooncup?
- Is it easy to use?

Jane Chiodini
Travel and Health Solutions

50

www.mooncup.co.uk

Jane Chiodini
Travel and Health Solutions

51

www.whizproducts.co/uk

Jane Chiodini
Travel and Health Solutions

52

Male travellers

20 – 29 year old age group at greater risk of accidents

Ref:McInnes R, Williamson L, Morrison A. (2002) Unintentional injury during foreign travel: a review. *Journal of Travel Medicine*, 9:297-307.

Jane Chiodini
Travel and Health Solutions

53

Medical History

"Cat Scan"

Jane Chiodini
Travel and Health Solutions

54

Medical History

1. Past and present medical history and current health status
2. Medication
3. Allergies to drugs or food/reaction to vaccination

Jane Chiodini
Travel and Health Solutions

55

Why consider the medical history ?

For example:


- Specialist advice may be needed e.g. those with severe renal or liver disease & malaria chemoprophylaxis
- Recent surgery or long term medical problems such as respiratory disease may impact of travel and fitness to fly
- Immunosuppression – some live vaccines contra-indicated, other vaccines may be less effective
- Impact on travel insurance with many medical problems
- Elderly people on regular medication need to be aware of continuing regular administration
- Establishing true anaphylaxis

Royal College of Nursing (2018) Competencies: Travel health nursing - career and competence development, London: RCN. Jane Chiodini
Travel health nurse



56

Implications of the questions we ask

Many sources to increase your knowledge and understanding of pre-travel risk assessment in more detail

The Yellow Book has a lot of information about medical history as do the National history databases



Jane Chiodini
Travel health nurse

57

Fitness to fly

The International Air Transport Association

- Is a trade association of the world's airlines. IATA supports airline activity and helps formulate policy and standards
- Its key priority is one of safety

<http://www.iata.org/publications/Documents/medical-manual.pdf>
<https://www.iata.org/publications/Pages/medical-manual.aspx>
Jane Chiodini
Travel health nurse

58



Aircraft Operations

Diagnosis	Assessment by a doctor with aviation medicine experience	Accept	Comments
CNS disorders (Central Nervous System)			
TIA (Stroke)	4 days or less	24 hours if stable or improving, with a normal ECG. Passenger travelling to the flight is under your advice should receive appropriate oxygen.	If an uncomplicated recovery has been made, further action is not required.
Cranial trauma	28 days or less	24 hours if generally well controlled.	
Cranial surgery	9 days or less	10 days, without loss of an and adequate general health.	
Cognitive impairment, Dementia	History of abnormal, unprovoked, aggressive or disturbed behaviour, discrimination, agitation or similar neurobehavioural syndrome; significant anxiety	100% impairment, independent function and living in the community, no significant paranoid, aggressive behaviour, hallucinations, or agitation. No change in medication since recent flight.	Consider support of travel companion
Gastro-intestinal			
Diagnosis	Assessment by a doctor with aviation medicine experience	Accept	Comments
GI bleed	28 hours or less following a bleed	2-10 days	10 days can travel if endoscopic or other clear evidence is felt has contributed to risk to indicate bleeding has ceased or healing. See also anemia.
Major abdominal surgery	9 days or less	2-10 days if uncomplicated recovery	e.g. bowel resection, open hysterectomy, renal surgery etc.
Acute abdomen	4 days or less	2-8 days if uncomplicated recovery	e.g. cholecystectomy (gall bladder removal), renal surgery.
Laparoscopic surgery	4 days or less	2-8 days if uncomplicated recovery	
Intestinal resection	28 hours or less	24 hours if gas absorbed	

http://www.iata.org/publications/Documents/medical-manual.pdf Jane Chiodini
Travel health nurse

59


Article – link on your page

Jane Chiodini
Travel health nurse

60

Travellers visiting friends and relatives (VFRs)




- Less likely to observe malaria chemoprophylaxis compliance
- Values and beliefs need to be explored

Chiodini PL, Patel D, Whitty CIM and Laloo DG. Guidelines for malaria prevention in travellers from the United Kingdom, London: Public Health England, November 2018. Jane Chiodini
Travel health nurse

61

Previous vaccine history



- In the absence of documentation, don't assume
- Ensure primary immunisations are up to date
- Give traveller a record of vaccines given

Jane Chiodini
Travel and Health Specialist

62

Destination


location, altitude, climate



Jane Chiodini
Travel and Health Specialist

63


Destination - political and economic situation



Jane Chiodini
Travel and Health Specialist

64

Departure date – season and timing




- Wet season – increases malaria risk
- Dry season – increases meningitis risk
- Last minute, still consider some vaccines e.g. hepatitis A

Jane Chiodini
Travel and Health Specialist

65

Length of stay

A 3 month visit carries a malaria risk around 6 times greater than a 2 week visit*




Complacency can creep in during a longer visit

* Chiodini PJ, Patel D, Whitty CJM and LaLoe DG - Guidelines for malaria prevention in travellers from the United Kingdom. London: Public Health England; November 2018

Jane Chiodini
Travel and Health Specialist

66


Mode of transport



Jane Chiodini
Travel and Health Specialist

67


Risk of accidents



Jane Chiodini
Travel nurse consultant

68

Purpose of trip and planned activities



People often seek adventure and take risks abroad they wouldn't consider when 'back home'.
People vary in their perception of risk

Jane Chiodini
Travel nurse consultant

69

Quality of accommodation




Top quality accommodation is not absolute assurance that there is no risk

Jane Chiodini
Travel nurse consultant

70

Financial budget



Those travelling on a tight budget may be at higher risk

General advice is not to eat at roadside stalls, however sometimes seeing the cooking process, quality of food and heat used, this may be a preferable option to a back street café!

Jane Chiodini
Travel nurse consultant

71

Health standards at the destination




- What are the healthcare standards like?
- The reuse of needles and syringes can be a common practice in some resource poor countries
- Carry a sterile medical kit
- Has adequate insurance been purchased?

Jane Chiodini
Travel nurse consultant

72

Collected the risk assessment information – then what?



Jane Chiodini
Travel nurse consultant

73

Do you know which resources we would use to make decisions?

WHERE TO GO

Jane Chiodini
Travel Health Specialist Nurse

74

Be aware of key UK resources for guidance

Jane Chiodini
Travel Health Specialist Nurse

75

UK National databases - www.travax.nhs.uk and for the public - www.fitfortravel.nhs.uk

Jane Chiodini
Travel Health Specialist Nurse

76

<https://travelhealthpro.org.uk/>

From NaTHNaC for healthcare professionals and the general public

Jane Chiodini
Travel Health Specialist Nurse

77

Access via your page and the 'New to Travel' page on my website

Please note, if using TRAVAX, you will still need a user name and password

<http://www.janechiodini.co.uk/tools/new-to-travel/>

Jane Chiodini
Travel Health Specialist Nurse

78

- Taking into consideration any patient specific factors (e.g. medical history, how high risk the destination is etc.) review the vaccines advised and decide what is needed – based also on previous vaccine history
- If a malarious area, also decide risk and identify appropriate chemoprophylaxis
- Consider advice required to manage the risks identified

Review the country specific information on a national database e.g. TravelHealthPro or TRAVAX

Jane Chiodini
Travel Health Specialist Nurse

79

Communicating the risk and providing advice

- Providing information about vaccines sufficient to provide adequate information to obtain informed consent
- Discussion of what is necessary and desirable – taking time and cost into the equation
- Advising on malaria prevention advice and deciding with patient the most suitable chemoprophylaxis
- Delivering other appropriate travel health advice – some will need to be in written format

Jane Chiodini
Travel Health Specialist Nurse

80

Advice leaflet that can be adapted for your use –



See item no. 4 at <http://www.janechiodini.co.uk/tools/> - written in Word format for you to adapt

Jane Chiodini
Travel Health Specialist Nurse

81

Food, water and personal hygiene advice

- Always wash hands before eating or preparing food
- Boiled water, bottled water - this includes ice cubes in drinks and water for cleaning your teeth
- Only eat well cooked fresh food
- Avoid leftovers and reheated food
- Ensure meat is thoroughly cooked
- Eat cooked vegetables, avoid salads
- Only eat fruit you can peel
- Never drink unpasteurised milk and avoid ice cream
- Shellfish is a high risk food

Jane Chiodini
Travel Health Specialist Nurse

82

- Studies have shown that the “Cook it, peel it, boil it or forget it” directive is not followed by many travellers and that conflicting results have been shown in the value of such strict advice
- New thinking in travel medicine is that food and drink can be placed into three categories
 - Safe
 - Probably safe
 - Unsafe
- There is no vaccine available for travellers’ diarrhoea

Ericsson CD. Prevention of Travelers Diarrhea in: Keystone J, Freedman D, Kazansky P, Connor B and Nordhult H. Eds. *Travel Medicine* 3rd Edition. Saunders, an imprint of Elsevier Inc; 2013. p. 191-196

Jane Chiodini
Travel Health Specialist Nurse

83

Food and beverage recommendations for travellers (this poster is on your page)

Category	SAFE	PROBABLY SAFE	UNSAFE
Beverages	<ul style="list-style-type: none"> • Carbonated soft drinks • Carbonated water • Boiled water • Purified water (iodine or chlorine) 	<ul style="list-style-type: none"> • Fresh citrus juices • Bottled water • Packaged (machine-made) ice 	<ul style="list-style-type: none"> • Tap water • Chipped ice • Unpasteurized milk
Food	<ul style="list-style-type: none"> • Hot, thoroughly grilled, boiled • Processed and packaged • Cooked vegetables and peeled fruits 	<ul style="list-style-type: none"> • Dry items • Hyperosmolar items (such as jam and syrup) • Washed vegetables and fruits 	<ul style="list-style-type: none"> • Salads • Sauces and 'salsa' • Uncooked seafood • Raw or poorly cooked meats • Unpeeled fruits • Unpasteurized dairy products • Cold desserts
Setting	Recommended restaurants	Local homes	Street vendors

Ericsson CD. Prevention of Travelers Diarrhea in: *Travel Medicine* 4th Edition, Eds. Keystone et al. Elsevier 2019

Jane Chiodini
Travel Health Specialist Nurse

84

Food water and personal hygiene advice... (this poster is on your page)



85

Travellers' diarrhoea advice

- **High risk areas** include North Africa, sub-Saharan Africa, the Indian Subcontinent, S.E. Asia, South America, Mexico and the Middle East
- **Medium risk areas** include the northern Mediterranean, Canary Islands and the Caribbean Islands
- **Low risk areas** include North America, Western Europe and Australia

Management

- Rehydration
- Anti diarrhoeal tablets
- Standby emergency treatment could be an option for some

Contact medical help if the affected person has:-

- A temperature
- Blood in the diarrhoea
- Diarrhoea for more than 48 hours (or 24 hours in children)
- Becomes confused

Jane Chiodini
Travel and Health

86

Prevention advice for hepatitis B, C and HIV infection

- Only accept a blood transfusion when essential
- If travelling to a resource poor country, take a sterile medical kit
- Avoid high risk procedures e.g. ear and body piercing, tattooing & acupuncture
- Avoid casual sex, especially without using condoms

Jane Chiodini
Travel and Health

87

Malaria prevention advice - the ABCD rules !



Photo credit: James Gathany

More information on malaria on day 2 of this course

Jane Chiodini
Travel and Health

88

Rabies advice

1. Do not touch any animal, even dogs and cats
2. If you are licked on broken skin, scratched or bitten in a country which has rabies, wash the wound thoroughly with soap and running water for 15 minutes then apply antiseptic.
3. Seek medical advice IMMEDIATELY, even if you have been previously immunised.

More information on rabies on day 2 of this course

Jane Chiodini
Travel and Health

89


Prevention of accidents advice

- Avoid alcohol and food before swimming
- Never dive into water where the depth is uncertain
- Only swim in safe water, check currents, sharks, jellyfish etc.
- Avoid alcohol when driving, especially at night
- Avoid hiring motorcycles and mopeds
- If hiring a car, rent a large one if possible, ensure the tyres, brakes and seat belts are in good condition
- Use reliable taxi firms, know where emergency facilities are

Jane Chiodini
Travel and Health

90

Risk management and the importance of documentation



Jane Chiodini
Travel and Health

91

Risk management form helps to 'record' best practice within the travel consultation

Form can be found in 'Tools' – item no. 2 <http://www.janechiodini.co.uk/tools/> Jane Chiodini
 Travel nurse & nurse prescriber

92

FOR HEALTH PROFESSIONAL USE ONLY IN CONJUNCTION with TRAVEL RISK ASSESSMENT FORM A

Patient Name: _____ dob: _____

Childhood immunisation history checked: _____ Additional information: _____

National database consulted for travel vaccines recommended for this trip and malaria chemoprophylaxis (if required): NaTHNaC: _____ TRAVAX: _____ Other: _____

Disease protection	Yes	Disease protection	Yes	Malaria Chemoprophylaxis Recommendation	Yes
BCG/Mantoux		Influenza		Atovaquone/proguanil	
Cholera		Meningitis ACWY		Chloroquine only	
Dip/tetanus/polio		MMR		Chloroquine and proguanil	
Hepatitis A		Rabies		Doxycycline	
Hepatitis B		TBE		Mefloquine	
Hepatitis A+B		Typhoid		Proguanil only	
Hepatitis A + Typhoid		Yellow fever		Emergency standby	
Japanese Encephalitis		Other		Weight of child:	

Vaccine and General Travel Advice required/provided

Potential side effects of vaccines discussed (including giving copy of the Patient Information Leaflet (PIL) from the vaccine packaging or obtainable from www.medicines.org.uk/emc/)

Patient consent for vaccination obtained: verbal written

Post vaccination advice given: verbal written

Jane Chiodini
 Travel nurse & nurse prescriber

93

Advice leaflet with additional resources – sample leaflet available on my website

See item no. 4 at <http://www.janechiodini.co.uk/tools/> - written in Word format for you to adapt Jane Chiodini
 Travel nurse & nurse prescriber

94

General travel advice leaflet given (all topics below in the surgery advice leaflet) and patient asked to read entire leaflet due to insufficient time to advise verbally on every topic: Yes / No

Items ticked below indicate topics discussed specifically within the consultation:

Prevention of accidents	Mosquito bite prevention	
Personal safety and security	Malaria prevention advice	
Food and water borne risks	Medical preparation	
Travellers' diarrhoea advice	Sun and heat advice	
Sexual health & blood borne virus risk	Journey/transport advice	
Rabies specific advice	Insurance advice	

Other specific specialised advice / information given on:
 e.g. smoking advice for a long haul flight; altitude advice; prevention of schistosomiasis, etc.

Source of advice used for further information: NaTHNaC TRAVAX Other

Jane Chiodini
 Travel nurse & nurse prescriber

95

Additional patient management or advice taken following risk assessment – for example

- Vaccine(s) patient declined following recommendation, and reason why
- Telephoned NaTHNaC or TRAVAX for advice or used Malaria Reference laboratory fax service
- Contacted hospital consultant for specific information in respect of a complex medical condition
- Identified specific nature/purpose of VFR travel

Jane Chiodini
 Travel nurse & nurse prescriber

96

Authorisation for a Patient Specific Direction (PSD)

Following the completion of a travel risk assessment, the below named vaccines may be administered under this PSD to:

Name: _____ dob: _____

Name, form & strength of medicine (generic/brand name as appropriate)	Dose, schedule and route of administration	Start and finish dates

Signature of Prescriber _____ Date _____

Jane Chiodini
 Travel nurse & nurse prescriber

97

- The template of this form could be adapted to use within a computer system, e.g. EMIS or Systm One
- If using paper copy of the form, then scan in after completion

Post Vaccination administration	
Vaccine details recorded on patient computer record (vaccine name, batch no., stage, site, etc.)	Y / N
SMS vaccines reminder or post card reminder service set up	Y / N
Travel record card supplied or updated:	Y / N
Travel risk management consultation performed by: (sign name and date)	Y / N

Form devised and created by Jane Chiodini © Updated May 2013

Jane Chiodini
Travel Health Consultant

98

Travel Health Training Ltd.
August 30 at 4:05 PM · G

I posted this image twice back in 2018 - the template has now moved on thanks to the great talent of a new nurse who the surgery who has improved its use and also made it SNOMED ready! To read more please go to a new blog I've written at <http://bit.ly/2ZEsvQZ> where you can also download it. And thank you so much for your help David Piercy - I really appreciate it!

EMIS WEB travel health template
Do you have one?

EMIS WEB SNOMED ready travel risk management template to download

Jane Chiodini's Blog
Entry: 30 August 2019
Saving time recording your travel consultation!

Back in the summer of 2018 I had an EMIS template that which allowed the use of my travel risk management form. Round at item no. 2 here. I put it out there for some of you to try and it generally received great feedback. The travel consultation is complex but sometimes writing up the information to provide evidence of all you covered and advised takes a long effort and significant time. However in my opinion this is essential but only as best practice but to also protect the practitioner.


<http://janechiodini.blogspot.com/2019/08/saving-time-recording-your-travel.html>

Jane Chiodini
Travel Health Consultant

99

Performing vaccination

Preparation of equipment and vaccines




Preparation of the patient

Jane Chiodini
Travel Health Consultant

100

Documenting the vaccinations

- Record of vaccines used must include the name of the drug, batch number, expiry date, site of administration and names of the administrator
- Ideally provide a written record of vaccinations given to the traveller



Variety of options now available, e.g. online, app format

Jane Chiodini
Travel Health Consultant

101


Conclusion

- No travel health consultation should take place without conducting a travel risk assessment and documenting all the information
- The assessment forms the basis of all subsequent decisions, advice given, vaccines administered and malaria prophylaxis advice that is offered
- Risk assessment and management takes time to perform correctly, and for best practice practitioners should leave sufficient time
- Good documentation is essential

Jane Chiodini
Travel Health Consultant

102

Case Study Practice



Travel health case studies
-
Putting theory into practice

Access via 'your page' to practice the case studies

Jane Chiodini
Travel Health Consultant

103




Agenda

- ✓ Introduction to travel medicine
- ✓ Travel risk assessment
- Travel vaccines and related issues
- Travel medicine operational issues
- Recap on resources

Jane Chiodini
Travel and Infectious Medicine


104

Vaccine preventable diseases and related issues



Jane Chiodini
Travel and Infectious Medicine

105




Agenda

- ✓ Travel vaccines and related issues
 - Key resources
 - Principles of vaccination and the rules
 - Range of vaccine preventable diseases and the specifics of these vaccines

Jane Chiodini
Travel and Infectious Medicine

106

Be aware of key UK resources for guidance




Yellow Book not online – may be in your workplace and some information is being placed on the NaTHNaC website

Jane Chiodini
Travel and Infectious Medicine

107

Key UK resources for guidance



JCVI (Joint committee on vaccination and immunisation)

Recommendations made by the JCVI to Government

Policy implemented and changes notified from Public Health England via direct communication

Information appears in VACCINE UPDATE along with other information re leaflets etc.

Jane Chiodini
Travel and Infectious Medicine

108

UK National databases - www.travax.nhs.uk and for the public - www.fitfortravel.nhs.uk



Jane Chiodini
Travel and Infectious Medicine

109

UK National databases - www.nathnac.org

Jane Chiodini
Thank you to Jane Chiodini

110

Working within National Guidelines and knowing one's limitations

Telephone helplines
NaTHNaC
 0845 602 6712
 Monday – Friday 9am to 11.00 am
 and 1pm to 2pm x 2
 Closed Monday and Friday at 2pm
 and other days at 3.30pm

TRAVAX
 0141 300 1130
 Mon. & Wed. 2 to 4pm
 Friday 9.30 to 11.30am

MRL e mail service
 Download risk assessment form
 from www.malaria-reference.co.uk, complete and e
 mail phe.malproph@nhs.net

E mail service – see malaria page, to be discussed next time

Jane Chiodini
Thank you to Jane Chiodini

111

Available from 'TOOLS' item no. 8

Jane Chiodini
Thank you to Jane Chiodini

112

Vaccines currently available to protect our travellers

Jane Chiodini
Thank you to Jane Chiodini

113

Diseases for consideration in this section?

PART 1
 NHS vaccines (mostly) and provided in an NHS setting (hepatitis B and meningitis can be private)

- Tetanus, diphtheria and polio
- Hepatitis A
- Typhoid
- Cholera
- Hepatitis B
- Meningitis

PART 2
 Always private, more specialist vaccines given by those more experienced

Covered on day 2

- Just touching on Yellow fever but separate training is required by NaTHNaC
- Rabies, Japanese B, tick-borne encephalitis

For more details regarding the charging of vaccines see FAQ no. 2 on my website
<http://www.janechiodini.co.uk/news/faqs/faq-no-2/>

Jane Chiodini
Thank you to Jane Chiodini

114

Immunisation timeline

Historical vaccine developments and introduction of vaccines in the UK

Historical information on NHS Choices – an interesting read see:
<http://www.nhs.uk/conditions/vaccinations/pages/the-history-of-vaccination.aspx>

Remember – this is item above accessed no. 14 to download from the 'Help' page at
<https://www.janechiodini.co.uk/help/immunisation-resources/>

Jane Chiodini
Thank you to Jane Chiodini

115

Other useful resources

Vaccination of individuals with uncertain or incomplete immunisation

A visual guide to vaccines used in the routine immunisation schedule

Jane Chiodini
Travel Health Specialist Nurse

116

UK and international immunisation schedules comparison tool

<https://www.gov.uk/government/publications/uk-and-international-immunisation-schedules-comparison-tool>

Jane Chiodini
Travel Health Specialist Nurse

117

Find them all via Immunisation resources in HELP

<https://www.janechiodini.co.uk/help/>

Jane Chiodini
Travel Health Specialist Nurse

118

Important to understand the principles of immunology

Jane Chiodini
Travel Health Specialist Nurse

119

Active Immunity

Jane Chiodini
Travel Health Specialist Nurse

120

Helpful videos for immunology and FAQs

<http://immunologyanimation.hpa.org.uk> and chapter 1 of the 'Green Book' access via the 'your dedicated page' on my website

Jane Chiodini
Travel Health Specialist Nurse

121

Another useful video published in May 2018

<https://www.ovg.ox.ac.uk/news/how-do-vaccines-work>

Jane Chiodini
Travavax & Travax

122

What factors might effect the immune response to a vaccine?

Age

Medical history

- Very young children (especially under 2 years) have difficulty developing an immune response to polysaccharide only vaccines, and conjugated vaccines are used where possible
- Immunocompromised individuals usually cannot receive live attenuated vaccines. Inactivated vaccines are usually safe, but their immune response may be inadequate

Jane Chiodini
Travavax & Travax

123

Contraindication to vaccinate?

- In general, a vaccine is absolutely contraindicated if a person has a confirmed anaphylactic reaction to a previous dose of the vaccine or product contained in the vaccine
- Pregnant women present a special risk group where, if the disease exposure is considered high during travel, most vaccine can be offered, although caution should be used with live vaccines
- All centres administering vaccines must be adequately prepared to deal with anaphylaxis

Jane Chiodini
Travavax & Travax

124

Knowledge of the route we give vaccines and how soon they start to work is needed

- Most vaccines given by IM or SC route except BCG and oral vaccines (cholera and live typhoid)
- An active immune response to vaccines begins within a few days of administration and peaks in approximately 10-14 days
- Primary vaccine courses need 2 or 3 doses to complete the series

Jane Chiodini
Travavax & Travax

125

Vaccine	Time until effective
BCG	6 weeks
Diphtheria	1-2 weeks after 3 rd dose
Hepatitis A (active)	2 weeks for optimum protection (the average incubation period for the disease is 28 days so it is often still useful to give the vaccine even at short notice prior to travel)
Hepatitis A immunoglobulin (passive)	Immediate
Hepatitis B	1 month after the 3 rd dose
Japanese encephalitis	7 days after the 2 nd dose, IXIARO® (Novartis) 1-2 weeks after the 2 nd dose, Green Cross vaccine (MASTA)
Measles/Mumps/Rubella (MMR)	2 weeks
Meningococcal vaccines (including ACW135Y)	2 weeks
Poliomyelitis (inactivated)	1-2 weeks after 3 doses
Rabies	1-2 weeks (after the 3 rd dose)
Tetanus	1-2 weeks after the 3 rd dose
Tick-borne encephalitis	2 weeks after the 2 nd dose
Typhoid injectable	1-2 weeks
Yellow fever	10 days

Time for Vaccines to become effective

Table adapted from TRAVAX

<http://www.travax.nhs.uk/vaccination/practice/arranging-schedules/time-for-vaccines-to-become-effective/>

Jane Chiodini
Travavax & Travax

126

The rules of vaccination

Jane Chiodini
Travel Health Specialist Nurse

General rules regarding vaccination

1. Administration of immunisation products is based on locally available immunisation schedules and the manufacturer's instructions. The manufacturer's instructions should be read and understood by the person administering the vaccine. The vaccine should be given to the person who is most likely to benefit from it.
2. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.
3. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.
4. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.
5. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.
6. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.
7. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.
8. If a vaccine is contraindicated, it should be avoided or postponed as far as possible. The person administering the vaccine should be aware of the reasons for this and should be able to explain them to the person receiving the vaccine.

on 'your dedicated' page

Jane Chiodini
Travavax & Travax


127

Peter thinks he has had 2 doses of hepatitis A vaccine in the past, but nothing is documented. He's going to travel to do some research work in a hospital in India – what would you do?

In the absence of documentation you cannot

ASSUME

the patient has been vaccinated, therefore further vaccines for protection should be given



Just2BChiodini
From: 10/10/19 to: 10/10/19


128

The evidence

For a variety of reasons, some individuals may not have been immunised or their immunisation history may be unknown.

If children and adults coming to the UK are not known to have been completely immunised, they should be assumed to be unimmunised and a full course of required immunisations should be planned.

Where a child born in the UK presents with an inadequate immunisation history, every effort should be made to clarify what immunisations they may have had. A child who has not completed the routine childhood programme should have the outstanding doses as described in the relevant chapters of the Green Book.



Page 6
REALLY IMPORTANT to read this chapter

https://www.gov.uk/government/publications/immunisation-schedule-the-green-book-chapter-11


Just2BChiodini
From: 10/10/19 to: 10/10/19

129

Michelle had a first hepatitis a vaccine at the age of one but never returned to complete the course, how would you proceed?

The evidence – chapter 11 again

Immunological memory from priming dose(s) are likely to be maintained in healthy individuals, increasing that interval will usually lead to a more pronounced response to the later dose. **Therefore, where any course of immunisation is interrupted, there is normally no need to start the course again - it should simply be resumed and completed as soon as possible.** Where vaccination was commenced some time previously however, the product received may have changed and the relevant chapter should therefore be consulted.



https://www.gov.uk/government/publications/immunisation-schedule-the-green-book-chapter-11

Just2BChiodini
From: 10/10/19 to: 10/10/19

130

The rules of vaccines

If a course goes off schedule and there is quite a long time interval, there is no need to re start the course, just pick up where it was left off and continue the course

ALWAYS USE AN AGE APPROPRIATE VACCINE

Department of Health. Immunisation against infectious disease (3rd Edition) London: TSO, 2006 Ch.17, p154
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/247564/Green-Book-Chapter-17.pdf


Just2BChiodini
From: 10/10/19 to: 10/10/19

131

Jon is backpacking around SE Asia for 6 months and is having a course of hepatitis B vaccine on a 0, 7 and 21 day schedule before he leaves. He had his day 0 dose today but won't be around for the day 7 dose and asks if he can attend in 5 days instead for his second dose. What would you advise?

The evidence

In general, it is acceptable to lengthen the intervals between doses and repeating previous vaccine doses is not necessary unless this is explicitly stated in the package insert. On the other hand, significant shortening of the intervals is not recommended



https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/247915/Green-Book-Chapter-4.pdf

Just2BChiodini
From: 10/10/19 to: 10/10/19

132

Site of injection and number that can be given at one time?

- If two or more injections need to be administered at the same time, they should be given in separate sites, preferably in a different limb. If more than one injection is to be given in the same limb, they should be administered at least 2.5cm apart
- Immunisations should not be given into the buttock, due to the risk of sciatic nerve damage and the possibility of injecting the vaccine into fat rather than muscle²




Figure 4.1 Preferred site for intramuscular and deep subcutaneous injections in older children and adults

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/247915/Green-Book-Chapter-4.pdf

2. Department of Health. Immunisation against infectious disease (3rd Edition) London: TSO, 2006 Ch.4, p29
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/247915/Green-Book-Chapter-4.pdf

Just2BChiodini
From: 10/10/19 to: 10/10/19

133

Where do you obtain relevant vaccine information?

- Green Book – for diseases and vaccines, online
- The National Databases (NaTHNaC and TRAVAX)
- Patient Group Directions - in your workplace
- Electronic Medicines Compendium - online
- British National Formulary – book or online
- Pharmaceutical companies - online

Jane Chiodini
Travax & NaTHNaC

134

Electronic Medicines Compendium www.medicines.org.uk

Jane Chiodini
Travax & NaTHNaC

135

Jane Chiodini
Travax & NaTHNaC

136

www.bnf.org

BNF Publications

Authoritative and practical.

Jane Chiodini
Travax & NaTHNaC

137

BNF and BNFC apps are great!

Jane Chiodini
Travax & NaTHNaC

138

Document your learnings

Disease prevention	Caustive organism	Mode of transmission prevention advice required	No. of vaccines in the course	Length of protection
Typhus, parvovirus and diphtheria				
Hepatitis A				
Typhoid				
Cholera				
Hepatitis B				
Meningococcal Meningitis				
Yellow fever				
Rabies				
Japanese encephalitis				
Tick borne encephalitis				

Jane Chiodini
Travax & NaTHNaC

139

Tetanus, polio and diphtheria

Jane Chiodini
Thank you to the NHS

140

Which disease is which?

Jane Chiodini
Thank you to the NHS

141

Learning more from Green Book chapters, but also www.nhs.uk

Disease	Organism	Mode of transmission
Tetanus	toxoid	Spores in the environment
Diphtheria	bacterium	Droplet infection
Polio	virus	Faecal oral and saliva

Jane Chiodini
Thank you to the NHS

142

but also www.nhs.uk

Jane Chiodini
Thank you to the NHS

143

Tetanus, diphtheria and inactivated polio vaccine in the immunisation schedule

Schedule above of 5 doses make the primary course in the UK.
Booster given if any of the three diseases are a risk at the destination, then an additional dose is provided on the NHS in a NHS setting.

Jane Chiodini
Thank you to the NHS

144

The routine immunisation schedule Route Autumn 2019

Tetanus, polio and diphtheria disease protection is not available in monovalent vaccines, only as one combined vaccine but in different products within the routine immunisation schedule

Infanrix hexa	DTaP/IPV/Hib/Hep B
Infanrix IPV	(DTaP/IPV)
Repevax	(DTaP/IPV)
Revaxis	(Td/IPV)

Jane Chiodini
Thank you to the NHS

145

Acronyms for vaccines and values of diphtheria content – D and d

Available vaccines^[6]

Diphtheria vaccines are available in two strengths according to dose of toxoid:

- High-dose - vaccines contain ≥ 30 IU of diphtheria toxoid and are used to achieve satisfactory primary immunisation of children - as in diphtheria/tetanus/acellular pertussis (DTaP) vaccine (capital D = high-dose).
- Low-dose - vaccines contain approximately 2 IU of toxoid and are used for primary immunisation of those aged over 10 years and for subsequent boosters (lower case d signifies low-dose as in dTaP).


https://www.cdc.gov/nczod/cv/ac/committees/quadrant/vac_advice.html
https://www.cdc.gov/nczod/cv/ac/committees/quadrant/vac_advice.html
https://www.cdc.gov/nczod/cv/ac/committees/quadrant/vac_advice.html

Jane Chiodini
Travel Health Specialist Nurse

146

Would this traveller need Td/IPV?

- Lucy is 19 years old and is going on a two week holiday in the Galapagos Islands
- She is up to date on all her scheduled national programme immunisations
- She hasn't travelled abroad before
- No PMH, she is on the OCP only



Jane Chiodini
Travel Health Specialist Nurse

147

Galapagos Islands off coast of Ecuador




Jane Chiodini
Travel Health Specialist Nurse

148

Would this traveller need Td/IPV?


- James is 26 years old
- he's taking a one year career break back packing around the world
- He last had a tetanus vaccine as a school booster 9½ years ago at the age of 16



Jane Chiodini
Travel Health Specialist Nurse

149

Keeping an eye on other groups



dTaP/IPV vaccine given between gestational weeks 20* and 32 rather than from week 28

*Can be given from 16 weeks but usually offered after the anomaly scan

Examples are ?

Boostrix-IPV or Repevax


Some very helpful resources on reverse of this chart

To find out more about whole cell pertussis and acellular pertussis see <http://www.who.int/biologicals/vaccines/pertussis/en/>

Jane Chiodini
Travel Health Specialist Nurse

150

FAQ on tetanus



Jane Chiodini
Travel Health Specialist Nurse

151

Public Health England
Preventing and improving the health of the nation

Tetanus
Guidance on the management of suspected tetanus cases and on the assessment and management of tetanus-prone wounds

Published November 2018

Chapter 30 Tetanus
November 2018

30 Tetanus

The disease

Tetanus is a wide disease caused by the action of tetanus toxin, released following infection by the bacterium Clostridium tetani. It is a life-threatening condition that may go undiagnosed. Tetanus is a disease of the body's soft tissue. The bacteria enter the body through a wound or scratch, or through a break in the skin.

The disease is characterised by generalised rigidity and spasms of skeletal muscles. The rigidity can occur in the neck, jaw and back, and is often associated with a 'lockjaw'.

Tetanus can occur at any age, but is most common in children under 15 years of age. It is a highly infectious and fatal disease, with a mortality rate of up to 100% if not treated.

Tetanus can occur in individuals because the disease is commonly present in the environment. It is a highly infectious and fatal disease, with a mortality rate of up to 100% if not treated.

History and epidemiology of the disease

Tetanus is a disease that has been known since the ancient Greeks in 1500 BC. It was first described in a medical text in 1812 by the German physician Johann Peter Frank. It was first described in a medical text in 1812 by the German physician Johann Peter Frank.

Between 1984 and 2017, there were 205 cases of tetanus reported from the UK. The majority of cases were reported in children under 15 years of age. The majority of cases were reported in children under 15 years of age.

Notes for public health practitioners

2018 and 2019

<https://www.gov.uk/government/publications/tetanus-immunoglobulin-recommendations-on-treatment-and-prophylaxis>

Jane Chiodini
Tetanus & Clostridium

152

Read the guidance to see definitions

6.2 Management of Tetanus-Prone Wounds

Tetanus-prone wounds* include:

- puncture-type injuries acquired in a contaminated environment and likely therefore to contain tetanus spores* e.g. gardening injuries
- wounds containing foreign bodies such as wood splinters**
- compound fractures
- wounds or burns with systemic sepsis
- certain animal bites and scratches**

*Note: individual risk assessment is required and this list is not exhaustive e.g. a puncture wound from discarded needle found in a park may be a tetanus-prone injury but a needlestick injury in a medical environment is not. **Similarly, although smaller bites from domestic pets are generally puncture injuries, animal saliva should not contain tetanus spores unless the animal has been rooting in soil or lives in an agricultural setting.

High-risk tetanus-prone wounds include:

- Any of the above with either:
 - heavy contamination with material likely to contain tetanus spores e.g. soil, manure
 - wounds or burns that show extensive devitalised tissue
 - wounds or burns that require surgical intervention that hours are high risk even if the contamination was not it

Includes UK born after 1961 with history of accepting vaccines

Immediate reinforcing dose of vaccine

Immediate reinforcing dose of vaccine

And on page 9 of the Green Book tetanus chapter

6.2 Management of Tetanus-Prone Wounds

Tetanus-prone wounds* include:

- puncture-type injuries acquired in a contaminated environment and likely therefore to contain tetanus spores* e.g. gardening injuries
- wounds containing foreign bodies such as wood splinters**
- compound fractures
- wounds or burns with systemic sepsis
- certain animal bites and scratches**

*Note: individual risk assessment is required and this list is not exhaustive e.g. a puncture wound from discarded needle found in a park may be a tetanus-prone injury but a needlestick injury in a medical environment is not. **Similarly, although smaller bites from domestic pets are generally puncture injuries, animal saliva should not contain tetanus spores unless the animal has been rooting in soil or lives in an agricultural setting.

High-risk tetanus-prone wounds include:

- Any of the above with either:
 - heavy contamination with material likely to contain tetanus spores e.g. soil, manure
 - wounds or burns that show extensive devitalised tissue
 - wounds or burns that require surgical intervention that hours are high risk even if the contamination was not it

Includes UK born after 1961 with history of accepting vaccines

Immediate reinforcing dose of vaccine

Immediate reinforcing dose of vaccine

And on page 9 of the Green Book tetanus chapter

Jane Chiodini
Tetanus & Clostridium

153

Jane Chiodini's Blog

Tuesday, 30 April 2019

Tetanus vaccine: 5 doses?

This isn't travel but because we have to also ensure patients are fully protected with the national immunisation schedule for tetanus I still think it's important.

A little history

For many years I've said once you have had 5 doses of tetanus containing vaccine then you are protected for life within the UK. Information written in the Green Book chapter on tetanus published in 2005 and 2007 stated that if you then sustained a tetanus-prone injury you would require treatment which would be tetanus-specific immunoglobulin but such documents are not available online any more.

The Patient info website on their page here states:

The primary course of three injections gives good protection for a number of years. The fourth and fifth doses (boosters) maintain protection. After the fifth dose, immunity remains for life and you do not need any further boosters (apart from some travel situations).

Notes for public health practitioners

2018 and 2019

Jane Chiodini
Tetanus & Clostridium

154

TRAVEL HEALTH PRO

COUNTRY INFORMATION LATEST NEWS OUTBREAK SURVEILLANCE IN UK

Home / Facebook / Poliomyelitis

POLIOMYELITIS

Poliomyelitis (polio) is a potentially paralyzing viral infection that only occurs in humans.

Content Sections

- Overview
- Risk areas
- Risk for travellers
- Transmission
- Signs and symptoms
- Diagnosis and treatment
- Preventing polio
- Vaccine information
- Resources

Recent and ongoing Polio Problems more about this on day 2

Jane Chiodini
Tetanus & Clostridium

155

Hepatitis A

Jane Chiodini
Tetanus & Clostridium

156

Hepatitis A

Hepatitis A, countries or areas at risk

The boundaries and names shown on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the validity or accuracy of its data, or derived from its information. Countries and territories are labeled as they appear on the most recent 1:50,000 scale maps available from the WHO Library of Reference Maps.


World Health Organization
11, rue de la Charité, 13300 Geneva, Switzerland
Tel: +41 (0)22 791 2111 Fax: +41 (0)22 791 3111
www.who.int

http://gamapserv.who.int/maplibrary/Files/Maps/Global_HepA_THRiskMap.png?ua=1

Jane Chiodini
Tetanus & Clostridium

157

Hepatitis A disease



<http://phli.cdc.gov/phli/home.asp>


- Viral infection
- Transmitted via contaminated food and water
- Those at higher risk – VFRs, long term travellers, those exposed to conditions of poor sanitation
- Incubation averages 28 - 30 days (range 15 to 50 days)
- Often asymptomatic in young children
- Abrupt onset of malaise, anorexia, nausea, fever followed by jaundice
- Fulminant hepatitis is more likely in those with pre-existing liver disease and in older individuals
- The overall case fatality ratio is low but is greater in older patients and those with pre-existing liver disease

<https://www.gov.uk/government/publications/hepatitis-a-the-green-book-chapter-17>

Jane Chiodini
Travellers' Clinician

158

Hepatitis A schedule – what the Green Book says



- The duration of protection from a completed course of vaccine can be expected to be at least 25 years and probably indefinite.
- However, PHE recommend that until further evidence is available on persistence of protective immunity, a booster dose at 25 years is indicated for those at ongoing risk of hepatitis A.

Hepatitis A vaccines can be used interchangeably: Chapter 4, page 145 of the Green Book

<https://www.gov.uk/government/publications/hepatitis-a-the-green-book-chapter-17>

Jane Chiodini
Travellers' Clinician

159

Five hepatitis A monovalent vaccines

VACCINE NAME	AGE GROUPS	WHEN TO BOOST- general principles and comments
HEPATITIS A vaccine (and schedules)		
• Hep A vaccine SPCs have different timings but note Ch. 4, 1 st paragraph in GB. Ideally, follow the summary of product characteristics but in late-presenting travellers, a course does not need to be restarted (DH 2013). Protection is expected for 25 years from the second dose – also see NaTHNaC info on Hepatitis A ** and detail below within the 'Key' section regarding GSK 'Havrix' vaccines***.		
VAQTA® Paediatric	1 - 17 years	Ideally, follow the summary of product characteristics but in late-presenting travellers, a course does not need to be restarted (DH 2013). Protection is expected for 25 years from the second dose – also see NaTHNaC info on Hepatitis A ** and detail below within the 'Key' section regarding GSK 'Havrix' vaccines***.
VAQTA® Adult	18 years and over	
Avaxim®	16 years and over	
Havrix Junior Monodose®	1 - 15 years	
Havrix Monodose®	16 years and over	

KEY
 * Within the Summary of Product Characteristics (SPC)
 ** The Green Book (2013) refers to all hep A products, so the 25 year protection also applies to the combined products and paediatric hepatitis A vaccines. Until further evidence is available on persistence of protective immunity, a further booster at 25 years is indicated for those at ongoing risk. See the Green Book chapter (page 154) and NaTHNaC document at www.nathnac.org/pro/factsheets/hep_a.htm
 *** SPC for Havrix Monodose & Havrix Junior Monodose April 2012 states 'Current recommendations do not support the need for further booster vaccination among immunocompetent subjects after 2 dose course'

See item no. 3 at <http://www.janechiodini.co.uk/tools/>

Jane Chiodini
Travellers' Clinician

160

Over to you!

1. What is the ideal schedule for a course of hepatitis A vaccine?
2. What do you do if the patient doesn't return for their hep A booster on time?
3. How long does a completed course of hepatitis A vaccine last?
4. Is the time of the protection taken from the first dose or booster dose of hepatitis A vaccine?
5. Are hepatitis A vaccines interchangeable?
6. If you gave a child a first hepatitis A vaccine and they return as an adult and a booster is required – which vaccine is best?
7. If a patient has a past medical history of confirmed hepatitis A infection, would you need to vaccinate them?
8. Could you give hepatitis A vaccine on the day of departure of a trip?
9. If a patient had HNIG recorded in their notes would you proceed on hep A protection?

Jane Chiodini
Travellers' Clinician

161

Over to you!

1. What is the ideal schedule for a course of hepatitis A vaccine? **0 and 6-12 months**
2. What do you do if they just boost when they turn up? **Just boost when they turn up**
3. How long does a completed course of hepatitis A vaccine last? **25 years**
4. Is the time of the protection taken from the first dose or booster dose of hepatitis A vaccine? **25 years from the booster dose**
5. Are hepatitis A vaccines interchangeable? **Yes**
6. If you gave a child a first hepatitis A vaccine and they return as an adult and a booster is required – which vaccine is best? **Boosting with an adult dose**
7. If a patient has a past medical history of confirmed hepatitis A infection, would you need to vaccinate them? **No**
8. Could you give hepatitis A vaccine on the day of departure of a trip? **Yes**
9. If a patient had HNIG recorded in their notes would you proceed on hep A protection? **Start a course of hep A vaccine**



<https://www.gov.uk/government/publications/hepatitis-a-the-green-book-chapter-17>

Jane Chiodini
Travellers' Clinician

162

Some historical hepatitis A protection/ vaccine information

Immunoglobulin given in 1980s and early 1990s but discontinued due risk of CJD from UK sourced blood products

Jane Chiodini
Travellers' Clinician

163

Some historical hepatitis A protection and vaccine information


- Immunoglobulin given in 1980s and early 1990s but discontinued due risk of CJD from UK sourced blood products
- Hepatitis A vaccine introduced in 1992 – Havrix – had to give two doses prior to travel as it had 720 ELISA units of hepatitis A (three doses in total course)
- Havrix Monodose available from 1994 which had 1440 ELISA units of hepatitis A and only one dose required prior to travel (two doses in total course)
- Vaqta Adult – problem in 1990s when some batches thought not to give protection – instructed at that time to disregard doses given previously and re-vaccinate. Vaqta Adult now available again
- See **Nuggets of Knowledge – hepatitis A**

Jane Chiodini
Travel Health Specialist Nurse

164

Would this traveller need hep A vaccine?


- Lucy is 19 years old and is going on a two week holiday in the Galapagos Islands
- She is up to date on all her scheduled national programme immunisations
- She hasn't travelled abroad before
- No PMH, she is on the OCP only
- Which vaccine schedule would you give?



Jane Chiodini
Travel Health Specialist Nurse

165

Would this traveller need hep A vaccine ?




- James is 26 years old
- he's taking a one year career break back packing around the world
- He last had a tetanus vaccine as a school booster 9½ years ago at the age of 16
- He tells you he had one dose of hep A vaccine when he was 12 years old but there is no record of it in the notes
- How would you proceed?

Jane Chiodini
Travel Health Specialist Nurse

166

Tip



Hepatitis A vaccine provides some of the most frequently asked questions therefore NaTHNaC and TRAVAX both have very helpful documents to help – it's a good idea to be aware of them

<http://travelhealthpro.org.uk/hepatitis-a/>

<http://www.travax.nhs.uk/diseases/vaccine-preventable/hepatitis-a/hepatitis-a-faqs.aspx>

FAQ on Hep A from TRAVAX

Jane Chiodini
Travel Health Specialist Nurse


167



Jane Chiodini
Travel Health Specialist Nurse


168

Recommend you do this to consolidate knowledge – see on your page



Jane Chiodini
Travel Health Specialist Nurse

169



170


WHO Factsheet – Hepatitis B updated July 2019

- Hepatitis B is a viral infection that attacks the liver and can cause both acute and chronic disease.
- The virus is transmitted through contact with the blood or other body fluids of an infected person.
- An estimated 257 million people are living with hepatitis B virus infection (defined as hepatitis B surface antigen positive).
- In 2015, hepatitis B resulted in 887 000 deaths, mostly from complications (including cirrhosis and hepatocellular carcinoma).
- Hepatitis B is an important occupational hazard for health workers.
- However, it can be prevented by currently available safe and effective vaccine.

<http://www.who.int/en/news-room/fact-sheets/detail/hepatitis-b>

Jane Chiodini
Thank you to CDC/NIH/USA

171




173

This Khmer woman died of hepatoma, four months after arriving in a refugee camp in Thailand

<http://www.vaccineinformation.org/hepb/photos.asp> or
<http://www.immunize.org/photos/hepatitis-b-photos.asp>

Jane Chiodini
Thank you to CDC/NIH/USA

174



175

USA and European resources: see 'immunisation resources' and 'hepatitis B' in 'HELP'

Jane Chiodini
Thank you to CDC/NIH/USA

176

Transmission of hepatitis B

Jane Chiodini
Thousand Oaks, CA, USA

177

A more recent craze – corset piercing

<http://unusual-things.blogspot.com/2011/05/surgeon-blasts-latest-craze-in-body.html>

Jane Chiodini
Thousand Oaks, CA, USA

178

The commonest transmission route of hepatitis B?

http://unicef-org.blogspot.co.uk/2010/10/unicef-executive-director-launches_31.html

Jane Chiodini
Thousand Oaks, CA, USA

179

Not travel related but important to understand

See page 14, chapter 18 of the Green Book

Age	Routine childhood programme	Babies born to hepatitis B infected mothers
Birth	1*	✓ Monovalent HepB
4 weeks	2	✓ Monovalent HepB
8 weeks	✓ DTaP/IPV/Hib/HepB	✓ DTaP/IPV/Hib/HepB
12 weeks	✓ DTaP/IPV/Hib/HepB	✓ DTaP/IPV/Hib/HepB
16 weeks	✓ DTaP/IPV/Hib/HepB	✓ DTaP/IPV/Hib/HepB
3 years of age	3*	✓ Monovalent HepB
3 years and 4 months	4*	✓ Test for HBsAg
		5*

* Give the recommended non-hepatitis B containing vaccines at per the routine schedule.

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/215622/gh_132617.pdf

<https://www.gov.uk/government/publications/hepatitis-b-antenatal-screening-and-newborn-immunisation-programme-best-practice-guidance>

Jane Chiodini
Thousand Oaks, CA, USA

180

Green Book recommendations for hepatitis B vaccine

- Injecting drug users
- Individuals who change sexual partners frequently, particularly MSM and commercial sex workers
- Close family contacts of a case or carrier
- Families adopting children from countries with a high or intermediate prevalence of hepatitis B
- Foster carers
- Individuals receiving regular blood or blood products and their carers
- Patients with chronic renal failure
- Patients with chronic liver disease
- Inmates of custodial institutions
- Individuals in residential accommodation for those with learning difficulties

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/148308/Green-Book-Chapter-18.pdf

Jane Chiodini
Thousand Oaks, CA, USA

181

The Green Book information regarding travellers

People travelling to or going to reside in areas of high or intermediate prevalence

Travellers to areas of high or intermediate prevalence who place themselves at risk when abroad should be offered immunisation. The behaviours that place them at risk will include sexual activity, injecting drug use, undertaking relief aid work and/or participating in contact sports. Travellers are also at risk of acquiring infection as a result of medical or dental procedures carried out in countries where unsafe therapeutic injections (e.g. the re-use of contaminated needles and syringes without sterilisation) are a risk factor for hepatitis B (Kane et al., 1999; Simonsen et al., 1999). Individuals at high risk of requiring medical or dental procedures in such countries should therefore be immunised, including:

- those who plan to remain in areas of high or intermediate prevalence for lengthy periods
- children and others who may require medical care while travelling to visit families or relatives in high or moderate-endemicity countries
- people with chronic medical conditions who may require hospitalisation while overseas e.g. dialysis
- those travelling for medical care

NB. The Green Book is nothing to do with whether the traveller should pay for vaccine or not

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/148308/Green-Book-Chapter-18.pdf

Jane Chiodini
Thousand Oaks, CA, USA

182

Schedules for hepatitis B vaccine

HEPATITIS B vaccine (and schedules)	Important – Hep B now in the childhood programme not included here
Engerix B® - 0, 1 and 6 months	Over 16 years
Engerix B® - 0, 1, 2 months	Over 16 years
Engerix B® - 0, 7, 21 days & 12 months	Over 18 years in SmPC – But also 16–18 years in Green Book
Engerix B® Paediatric 0, 1, 6 months	0 to 15 years
Engerix B® Paediatric 0, 1, 2 months	0 to 15 years
Engerix B® Option of two doses of 1 ml (20mcg) for low-compliance adolescents given 6 months apart when the risk of hepatitis B is low and completion of course can be assured before risk is high	11–15 years
HBVaxPRO® 0, 1, and 6 months	16 years and over
HBVaxPRO® 0, 1, 2 months	16 years and over
HBVaxPRO® Paediatric 0, 1 & 6 months	0–15 years
HBVaxPRO® Paediatric 0, 1, 2 months	0–15 years

Note: 0, 1, 2 month schedule Green Book doesn't advise 4th dose at 12 months unless they remain at continued high risk, see Ch.18 Page 16. Note SmPCs do advise a 4th dose but GB should be followed. Update to policy in the Green Book in June 2017 for hepatitis B for all (which would include travellers) states those who have received a primary course do not require a reinforcing dose of hep B containing vaccine except health care workers (boost once at 5 years), patients with renal failure and at time of significant exposure. Please read Ch. 18 page 13 of Green Book for detail. Testing for evidence of immunity post immunisation is not routinely recommended. See GB, Ch.18, Page 18

Two products, four presentations

Jane Chiodini
Travellers' Clinic UK Ltd

183

Which schedule?

Hepatitis B Green Book chapter page 12

Pre-exposure immunisation schedule for **high risk individuals**

- For pre-exposure prophylaxis in most adult and childhood risk groups, an accelerated schedule should be used, with vaccine given at zero, one and two months.
- Higher completion rates are achieved with the accelerated schedule (at zero, one and two months) in groups where compliance is difficult (e.g. in people who inject drugs [PWID] and genitourinary medicine clinic attenders) (Asboe et al., 1996).
- This improved compliance is likely to offset the slightly reduced immunogenicity when compared with the zero-, one- and six-month schedule, and similar response rates can be achieved by the opportunistic use of a fourth dose after 12 months.
- An alternative schedule at zero, one and six months should only be used where rapid protection is not required and there is a high likelihood of compliance.
- If the primary course is interrupted it should be resumed but not repeated.)

Jane Chiodini
Travellers' Clinic UK Ltd

184

What about hepatitis B boosters?

Hepatitis B Green Book chapter page 13

Reinforcing doses for those who have received pre-exposure immunisation

The current UK recommendation is that those who have received a primary course of immunisation, including children vaccinated according to the routine childhood schedule and individuals at high risk of exposure, **do not require a reinforcing dose of Hep B-containing vaccine**, except in the following categories:

- healthcare workers (including students and trainees), who should be offered a single booster dose of vaccine, once only, around five years after primary immunisation
- patients with renal failure
- at the time of a significant exposure (see the chapter for more detail)

Jane Chiodini
Travellers' Clinic UK Ltd

185

Public Health England
Protecting and improving the nation's health

Statement in this guidance Feb 2018

– not yet in the Green Book

Plan for phased re-introduction of hepatitis B vaccine for lower priority groups in 2018


Booster doses in healthcare workers

On the advice of the Joint Committee on Vaccination and Immunisation (JCVI), boosters (priority group 5) will no longer be routinely required in healthy, immunocompetent adults who have completed a primary course of vaccine, including healthcare workers who are known responders.

See page 8

Jane Chiodini
Travellers' Clinic UK Ltd

186



In summary for hepatitis B vaccine given for travel purposes

- Use 0, 1 and 2 month schedule in preference to 0, 1 and 6 month when more rapid protection is needed
- If insufficient time before travel, use a 0, 7, 21 day and then reinforce at 12 months
- No longer boost at 5 years for travel
- Blood test not routinely performed for seroprotection in travellers

Jane Chiodini
Travellers' Clinic UK Ltd

187

Where would you find information about the levels of protection?



emc®

Start searching the eMC here

Pharmacodynamic properties

Pharmacodynamic group: Hepatitis B vaccine, ATC code: J07DC01

Efficacy: Efficacy is defined as the percentage of subjects with anti-HBc antibody concentrations ≥ 10 IU/L observed in clinical studies with the different schedules mentioned in section 4.2.

Population	Schedule	Seroprotection rate
Healthy subjects (0 to 65 including 15 years of age)	0, 1, 6 months	at least 71.00%
	0, 1, 2, 12 months	at least 76.00%
	0, 1, 2, 12 months	at least 79.00%

Jane Chiodini
Travellers' Clinic UK Ltd

188

Hepatitis A vaccine antigen content within vaccines


Adapted from table 1 of the Public Health England Hepatitis A vaccination in adults temporary recommendations *

Hep A vaccine formulation	Trade name	Hep A vaccine antigen content	Adult dose Hep A antigen equivalent	Made by
Adult monovalent hep A	AVAXIM	160 U	Full dose	SP
	HAVRIX MONODOSE	1440 EU	Full dose	GSK
	VAQTA ADULT	50 U	Full dose	MSD
Paediatric monovalent hep A	HAVRIX JUNIOR MONODOSE	720 EU	Half dose	GSK
	VAQTA PEDIATRIC	25 U	Half dose	MSD
Adult combination hepatitis A/B	TWINRIX ADULT	720 EU	Half dose	GSK
Paediatric combination hepatitis A/B	TWINRIX PEDIATRIC	360 EU	Quarter dose	GSK
	AMBIRIX	720 EU	Half dose	GSK
Combination hepatitis A/typhoid	VIATIM	160 EU	Full dose	SP

KEY
SP – Sanofi Pasteur; GSK = GlaxoSmithKline; MSD = Merck Sharpe Dohme Ltd.
<https://www.gov.uk/government/publications/hepatitis-a-infection-prevention-and-control-guidance>
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/626228/hepatitis_a_vaccination_recommendations.pdf

Made into a poster and now on your webpage

189



191

Combination A and B vaccines schedules

- Twinrix adult**
 - 0, 1 & 6 months from 16 years
 - 0, 7, 21 days & 12 mths (18yrs)
- Twinrix paediatric**
 - 0, 1 & 6 months
 - Use in 1 – 15 year age group
- Ambirix**
 - 0 and 6 - 12 months
 - Use in 1 to 15 years

Jane Chiodini
The Health Education Unit

192

Hepatitis A vaccine antigen content within vaccines

Adapted from table 1 of the Public Health England Hepatitis A vaccination in adults temporary recommendations *

Hep A vaccine formulation	Trade name	Hep A vaccine antigen content	Adult dose Hep A antigen equivalent	Made by
Adult monovalent hep A	AVAXIM	160 U	Full dose	SP
	HAVRIX MONODOSE	1440 EU	Full dose	GSK
	VAQTA ADULT	50 U	Full dose	MSD
Paediatric monovalent hep A	HAVRIX JUNIOR MONODOSE	720 EU	Half dose	GSK
	VAQTA PEDIATRIC	25 U	Half dose	MSD
Adult combination hepatitis A/B	TWINRIX ADULT	720 EU	Half dose	GSK
Paediatric combination hepatitis A/B	TWINRIX PEDIATRIC	360 EU	Quarter dose	GSK
	AMBIRIX	720 EU	Half dose	GSK
Combination hepatitis A/typhoid	VIATIM	160 EU	Full dose	SP

KEY
SP – Sanofi Pasteur; GSK = GlaxoSmithKline; MSD = Merck Sharpe Dohme Ltd.
<https://www.gov.uk/government/publications/hepatitis-a-infection-prevention-and-control-guidance>
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/626228/hepatitis_a_vaccination_recommendations.pdf

Made into a poster and now on your webpage

193

Combination hepatitis B vaccines used in the UK – from the Green Book

Table 18.2 Dosage of combined hepatitis A and hepatitis B vaccines by age (from the Green Book)

Vaccine product	Ages	Dose HAV	Dose HBV	Volume
Twinrix Adult® Hepatitis A (inactivated) and hepatitis B (rDNA) (HAB) vaccine (adsorbed)	16 years or over	720 ELISA units	20µg	1.0ml
Twinrix Paediatric® Hepatitis A (inactivated) and hepatitis B (rDNA) (HAB) vaccine (adsorbed)	1 – 15 years	360 ELISA units	10µg	0.5ml
Ambirix® Hepatitis A (inactivated) and hepatitis B (rDNA) (HAB) vaccine (adsorbed)	1 – 15 years	720 ELISA units	20µg	1.0ml

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/148308/Green-Book-Chapter-18.pdf

Jane Chiodini
The Health Education Unit

195

Tip

In practice we give hepatitis B for lifestyle risks and travel, but not usually occupational risk. Hepatitis B is a large topic – it would be useful to read the Green Book chapter on this topic at https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/148308/Green-Book-Chapter-18.pdf


And the BMA guidance document at <http://bma.org.uk/practical-support-at-work/doctors-as-managers/managing-your-practice/focus-hepatitis-b-immunisations>

See the hepatitis B document on **your dedicated page !**

Jane Chiodini
The Health Education Unit

196

Typhoid




Jane Chiodini
Thailand 2018-2019

197

Enteric fevers – typhoid & paratyphoid

- Gram-negative bacterial infection
- Transmission by faecal-oral route, water borne and human to human
- Those at higher risk include VFRs, young children, long term travellers and those exposed to conditions of poor sanitation – mainly in Asia
- Incubation 7 to 14 days
- Fever, chills, headache, malaise, weakness, anorexia, abdominal pain, diarrhoea
- Complications in 10% -15%: intestinal perforation, bacteraemia, meningitis
- Chronic carrier status in <3% infected persons



Jane Chiodini
Thailand 2018-2019

198

Data from PHE

Public Health England
Enteric fever (typhoid and paratyphoid) England, Wales and Northern Ireland: 2017




Figure 3. Laboratory-confirmed cases of enteric fever, England, Wales and Northern Ireland by travel history: 2008 to 2017

■ Travel abroad not stated ■ No travel abroad
■ New entrant to UK ■ Foreign visitor to UK
■ Travelled abroad from UK — % Travel history
— % travel abroad


<https://www.gov.uk/government/publications/typhoid-and-paratyphoid-laboratory-confirmed-cases-in-england-wales-and-northern-ireland>

Jane Chiodini
Thailand 2018-2019

199

Reason for travel and destination

Figure 4. Reason for travel for laboratory-confirmed cases of enteric fever that travelled abroad from England, Wales and Northern Ireland: 2017 (N=287)



India, Pakistan and Bangladesh were the highest risk country for travellers returning with enteric fever

Table 3. Countries of travel and ethnicity for laboratory-confirmed cases of enteric fever that travelled abroad from England, Wales and Northern Ireland to visit friends and relatives: 2017 (N=203)*

Presumed country of infection	Ethnicity						Total
	Indian	Pakistani	Bangladeshi	Asian other	Black African	Other/mixed	
India	84	1	-	-	-	2	10
Pakistan	1	55	1	1	-	-	8
Bangladesh	-	-	20	-	-	-	1
Sub-Saharan Africa	-	-	-	-	5	-	1
Other Asia	1	-	-	4	2	5	1
Total	86	56	21	5	7	7	21


<https://www.gov.uk/government/publications/typhoid-and-paratyphoid-laboratory-confirmed-cases-in-england-wales-and-northern-ireland>

Jane Chiodini
Thailand 2018-2019

200

Two vaccines now for typhoid protection

TYPHOID vaccine		
Typhim VI® Single dose	2 years and over	3 years
Vivotif® (Ty21a) Oral vaccine on days 0, 2 & 4	5 years and over	3 years (Take with cold or luke warm drink 1 hr before meal, swallow capsule whole)



Injectable typhoid protection is a polysaccharide vaccine and so just one dose makes up 'the course'. After this time period if further typhoid protection is needed a new dose is given.

Jane Chiodini
Thailand 2018-2019

201

Details from the Green Book

Dosage of injectable monovalent typhoid vaccines			
Vaccine product	Ages	Dose	Volume
Typhim VI	Two years and older**	25µg	0.5ml
Typherix (discontinued in 2018)	Two years and older*	25µg	0.5ml

Dosage of oral monovalent typhoid vaccine		
Vaccine product	Ages	Dose
Vivotif	Five years and older	Three capsules on days 0, 2 and 4

Dosage of combined typhoid and hepatitis A vaccines**			
Vaccine product	Ages	Dose typhoid	Dose HAV†
Hepatitrix (discontinued in 2018)	15 years and older	25µg	1440 EUISA units
VIATIM	16 years and older	25µg	160 antigen units

* Children between the ages of 12 months and two years should be immunised off-license if following a detailed risk assessment the risk of typhoid fever is considered high.
 ** For booster doses of either typhoid or HAV, single antigen vaccines can be used
 † HAV – hepatitis A vaccine

Note
 Typherix and Hepatitrix have now been discontinued – as now highlighted in the Green Book
 Children between the ages of 12 months and two years should be immunised off-license if following a details risk assessment the risk of typhoid fever is considered high

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/49512/Green_Book_Chapter_33_04_120318.pdf

Jane Chiodini
Thailand 2018-2019

202


Oral typhoid vaccine - Vivotif

- LIVE vaccine*
- Use from 5 years of age
- Three doses on days 0, 2 & 4 (Green Book)

Administration

- The capsule should be taken approximately one hour before a meal with a cold or lukewarm drink (temperature not to exceed body temperature, e.g. 37°C)
- The vaccine capsule should not be chewed and should be swallowed as soon as possible after placing in the mouth

* can be administered at any time before or after other live vaccines.



Jane Chiodini
Travel Nurse Consultant

203

Moved to Chapter 11 of the Green Book – in April 2019




Table 11.2 Recommended time intervals when giving more than one live attenuated vaccine

Vaccine combinations	Recommendations
Yellow fever and MMR	A four week minimum interval should be observed between the administration of these two vaccines. Yellow fever and MMR should not be administered on the same day.
Vivotif (and zoster vaccine and MMR)	If these vaccines are not administered on the same day, then a four week minimum interval should be observed between vaccines.
Tuberculin skin testing (Mantoux) and MMR	MMR vaccination and tuberculin skin testing can be performed on the same day. However, if a tuberculin skin test has already been initiated, then MMR should be delayed until the skin test has been read, unless protection against measles is required urgently. If a child has had a recent MMR, and requires a tuberculin test, then a four week interval should be observed.

All currently used live vaccines (BCG, rotavirus, live attenuated influenza vaccine (LAIV), oral typhoid vaccine, yellow fever vaccine, zoster and MMR).


Co-administration of these two vaccines can lead to sub-optimal antibody responses to yellow fever, mumps and rubella antigens (Nascimento et al, 2011). Where protection is required rapidly then the vaccines should be given at any interval; an additional dose of MMR should be considered.


Jane Chiodini
Travel Nurse Consultant

204

Oral typhoid - Vivotif®

www.medicines.org.uk





Jane Chiodini
Travel Nurse Consultant

205

Interactions with other medication from the SmPC

- Antibiotics – Vivotif may not work if it is taken while you are also taking antibiotics. Take Vivotif no earlier than 3 days after the last dose of an antibiotic
- Medicines to prevent malaria – do not start these until 3 days after the last dose of Vivotif
- Yellow fever vaccine can be given while taking Vivotif

Jane Chiodini
Travel Nurse Consultant

206

Charging and administration

Oral typhoid vaccine is an NHS provision in an NHS setting

1. You could buy this vaccine in and bring the patient in for all three doses
2. You could administer the vaccine to the patient for the first dose and given them the other two doses to take home to self administer but the vaccine must be stored at 2 – 8°C
3. You could supply the vaccine on an FP10 and allow the patient to self administer

You need to ensure that your traveller understands the importance of, and can assure the cold chain in points 2 and 3 above.

Jane Chiodini
Travel Nurse Consultant

207


Prioritising typhoid vaccine if there is a shortage – who would you give it to?

- Family of four going to an all inclusive break for a 10 days Cancun
- 40 year old couple travelling to stay in 4 star hotel in Bangkok for two weeks
- 26 year old man going to stay in Bangkok in a guest house/hostel type accommodation and he has type 1 diabetes
- Parents and their children of 7, 5 and 2 years travelling to Pakistan for 3 weeks to see family

Jane Chiodini
Travel Nurse Consultant

208

Tip



NaTHNaC and TRAVAX have both written information documents on typhoid as well

<http://travelhealthpro.org.uk/typhoid-and-paratyphoid/>

<http://www.travax.nhs.uk/diseases/vaccine-preventable/typhoid/typhoid-faqs.aspx>

Jane Chiodini
Travel Health Pro

209

Hepatitis A + Typhoid available as a combined vaccine

Why give combination vaccines?

COMBINED vaccines (and schedules)		
VIATIM® (Hepatitis A and typhoid) Single dose	16 years and over	6-12 months for hepatitis A component (then hepatitis A booster given as a monovalent vaccine) and 3 years for the typhoid component


The combination hepatitis A and typhoid vaccine can be given with the hep A protection provided as either the first dose or reinforcing or booster dose of hep A vaccine as long as there is the three year interval to fulfill the typhoid requirement of the vaccine.

Jane Chiodini
Travel Health Pro

210

Does this traveller need typhoid vaccine ?

Anu is travelling to Mumbai to see relatives for a 4 week stay – she is 22 years old. She had a 1st dose hepatitis A vaccine at the age of 14 years. Would she need a typhoid vaccine and which one would you give if so?



Anu needs a booster dose of hepatitis A vaccine and she needs a typhoid vaccine so it would be very appropriate in this situation to give her a combined hepatitis A and typhoid vaccine.

Jane Chiodini
Travel Health Pro

211

Food and water hygiene advice remain paramount





<https://travelhealthpro.org.uk/news/349/extensively-drug-resistant-typhoid-fever-in-pakistan>

Jane Chiodini
Travel Health Pro

212

Cholera






Jane Chiodini
Travel Health Pro

213

Cholera

Disease	Organism	Mode of transmission
Cholera	Bacterial infection	Mainly water-borne through ingestion of faecally contaminated water or shellfish and other foods. Person-to-person spread may occur through the faecal-oral route

<http://gamespinner.who.int/magislibrary/>

Jane Chiodini
Travel Health Pro

214

Cholera is referred to in 'pandemics' – currently in the 7th Pandemic

Countries reporting cholera, 2010-2015

Countries reporting cholera, 2010-2014

Countries reporting cholera, 2015

The boundaries and names shown and the designation used in this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, or area or of its boundaries, or concerning the delimitation of its frontiers or boundaries. Cities and towns not shown are not necessarily shown in this map.

© 2015 World Health Organization

© 2015 World Health Organization

<http://gamapserver.who.int/maplibrary/>

Jane Chiodini
Thank you to Jane Chiodini

215

<http://globalhealth.unc.edu/blog/2011/06/haiti-diary-back-to-the-basics/>

Jane Chiodini
Thank you to Jane Chiodini

216

© Reuters

Haitian cholera victims receive treatment inside a hospital run by Doctors Without Borders in Port-au-Prince. The death toll from Haiti's cholera epidemic has reached more than 900

16 November 2010

<http://www.dailymail.co.uk/news/article-1330282/Anti-UN-riots-Haiti-leave-people-dead-locals-blame-cholera-outbreak-UN-peacekeepers.html>

Jane Chiodini
Thank you to Jane Chiodini

217

Cholera

- Acute intestinal infection
- Causal bacterium - *Vibrio cholerae*
- Transmitted faecal orally
- 90% cases are mild to moderate
- 10% cases very severe – leading to profuse diarrhoea, vomiting, circulatory collapse and shock
- Mortality rate can be over 50% in untreated cases, unless rapid rehydration therapy is given promptly
- Chronic carriage is rare
- Organism survives for up to 2 weeks in fresh water and 8 weeks in salt water
- Transmission normally through infected drinking water

Jane Chiodini
Thank you to Jane Chiodini

218

Management of cholera

- Fluid replacement
- Prompt action improves outcome
- IV fluids in severe cases or when vomiting
- Rapid rehydration until signs improve
- NG tube used if IV not possible
- Antibiotic therapy in severe cases

Jane Chiodini
Thank you to Jane Chiodini

219

Cholera recommendations for administration

Advice from the Green Book

- Immunisation against cholera can be considered, following a full risk assessment, for the following categories of traveller:
 - relief or disaster aid workers
 - persons with remote itineraries in areas where cholera epidemics are occurring and there is limited access to medical care
 - travellers to potential cholera risk areas, for whom vaccination is considered potentially beneficial.

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/263838/Green-Book-Chapter-4-v2_0.pdf

Jane Chiodini
Thank you to Jane Chiodini

220

Cholera vaccine fact finding


- What is the youngest age at which give cholera vaccine can be prescribed?
- How many doses would you give a child?
- How many doses would you give an adult?
- What is the minimum and maximum time interval between doses?
- How long does cholera vaccine last?

Jane Chiodini
Travel Nurse Consultant

221

Cholera Vaccine

CHOLERA vaccine	2 years and over	2 yrs in age 6 to adult 6 months in 2 – 6 year olds NBM 1 hr before & after vaccine
Dukoral® Oral vaccine. 2 doses, minimum 1 wk. apart and maximum 6 weeks apart, from 6yrs of age. 3 doses, in 2 – 6 year olds		



Food and drink should be avoided 1 hour before and 1 hour after vaccination. Oral administration of other medicinal products should be avoided within 1 hour before and 1 hour after administration of Dukoral.

Jane Chiodini
Travel Nurse Consultant

222

From the PiL


Instructions:




1. To prepare buffer solution dissolve the effervescent granules in a glass of cool water (approx. 150 ml). Do not use any other liquid.
Children 2-6 years: pour away half of the buffer solution.
2. Shake the vaccine bottle (1 bottle = 1 dose).
3. Add the vaccine to the buffer solution. Mix well and drink the mixture. Drink the vaccine within 2 hours after mixing with the buffer solution. Avoid food and drink starting 1 hour before until 1 hour after the vaccination.

Jane Chiodini
Travel Nurse Consultant


223

For further information to learn more about these diseases, look at the Green Book (online) NaTHNaC and TRAVAX



Tip



Jane Chiodini
Travel Nurse Consultant

225

Meningococcal meningitis



Jane Chiodini
Travel Nurse Consultant

226

Meningococcal Meningitis

- Meningococcal disease is a rare, but potentially devastating infection
- Caused by the bacteria *Neisseria meningitidis* of which there are 6 disease-causing strains called serogroups (A, B, C, W, Y and X)
- Approximately 10 percent of the general population of the UK are thought to carry *N. meningitidis* in the lining of the nose and throat
- Spread between individuals occurs through coughing, sneezing, kissing or during close contact with a carrier
- Carriers do not have symptoms, but can develop disease when bacteria invade the bloodstream from the nasopharynx
- Invasive disease is a rare but serious outcome usually presenting as septicaemia or meningitis


<http://travelhealthpro.org.uk/diseases/meningococcalmeningitis/>

Jane Chiodini
Travel Nurse Consultant

227

Meningococcal Meningitis

- Less commonly, individuals may present with pneumonia, myocarditis, endocarditis, pericarditis, arthritis, conjunctivitis, urethritis, pharyngitis and cervicitis




- The incubation period is from two to seven day

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/223749/Green_Book_Chapter_22_v2_3.pdf

Jane Chiodini
Travel and Vaccination

228

Meningococcal meningitis vaccine given to travellers going to meningitis belt in Africa



http://gamapserver.who.int/maplibrary/files/Maps/Global_MeningitisRisk_11thRiskMap.png
<http://www.who.int/travel/yellowbook/2014/infectious-diseases-related-to-travel/meningococcal-disease>

Jane Chiodini
Travel and Vaccination

229

Vaccine recommendation for travellers to the meningitis belt

Meningococcal disease vaccination
 Vaccination is recommended for those whose activities or medical condition put them at increased risk including:

- healthcare workers
- those visiting friends and relatives
- those who live or travel 'rough' such as backpackers
- long-stay travellers who have close contact with the local population
- those with certain rare immune system problems (complement disorders) and those who do not have a functioning spleen

From NaTHNaC ←

Who Should I Vaccinate for Meningococcal Meningitis?

Consider vaccinating:

- Travellers who are likely to have close, prolonged contact with the local population
- Long stay travellers
- Those visiting friends and relatives
- Those who will be exposed to crowded areas (e.g. stadia, schools, dormitories, hospitals)
- Travellers visiting an area affected by an ongoing outbreak or epidemic
- Immunocompromised travellers (including asplenia) visiting endemic areas

→ **From TRAVAX**

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/292994/Green_Book_Chapter_22_v2_5.pdf


Jane Chiodini
Travel and Vaccination

230

and travellers going on a pilgrimage to Umrah and Hajj

The KSA Ministry of Health (MoH) currently recommends that, as a precautionary measure, pregnant women and young children, should postpone the performance of the Hajj and Umrah.

Those with severe medical conditions such as terminal cancers, advanced cardiac, respiratory, liver, kidney diseases or senility are exempt from these religious duties



Write 5 years on the certificate for visa purposes now and it must specify the type of vaccine

Note – information given that vaccine must be given minimum of 10 days prior to entry into the country

Certificates available at:
https://hsc.gsk.co.uk/content/dam/global/hcpportal/en_GB/therapyaareas/vaccines/pdfs/meningococcal-acwy-certificate.pdf
 and <https://drivetravaccines.medica.com/medica?wicketinterface=3...>

Jane Chiodini
Travel and Vaccination

231

The ACWY vaccines

<p>Menveo</p> <ul style="list-style-type: none"> Conjugate vaccine Use from two years of age Available from GSK GSK data gives 5 years protection from administration 	<p>Nimenrix</p> <ul style="list-style-type: none"> Conjugate vaccine Use from 6 weeks now* Just had black triangle removed Available from Pfizer Pfizer studies up to 60 months – refer to pharmacodynamics properties in the SPC
--	---

* If needing to give, please check the Green Book, the SPC and TravelHealthPro


https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/292994/Green_Book_Chapter_22_v2_5.pdf

Jane Chiodini
Travel and Vaccination

232

When would you boost - it's been very confusing!

- The Joint Committee on Vaccination and Immunisation (JCVI) Committee reviewed information on length of protection following ACWY conjugate vaccination. Antibody against serogroup A disease was the first to wane, and this meant boosting was important for travel, but less important for the routine Men ACWY programme in the UK.
- For travellers at continued risk, the Committee agreed that boosting every five years would be a sensible approach until data became available.





<http://www.who.int/wer/2016/wer1626-27.pdf?ua=1>


Jane Chiodini
Travel and Vaccination

233

For further information to learn more about these diseases, look at the Green Book (online) NaTHNaC and TRAVAX

Tip



Jane Chiodini
Travel Health Specialist Nurse

234


Agenda

- ✓ Introduction to travel medicine
- ✓ Travel risk assessment
- ✓ Travel vaccines and related issues
 - Travel medicine operational issues
 - Recap on resources

Jane Chiodini
Travel Health Specialist Nurse

235

Immunisation Training




Jane Chiodini
Travel Health Specialist Nurse

236

Jane Chiodini
Travel Health Specialist Nurse


Some of the resources on the 'Help' page with more being added



Jane Chiodini
Travel Health Specialist Nurse


237

Jane Chiodini
Travel Health Specialist Nurse



IMMUNISATION RESOURCES


For example, Immunisation Resources



Jane Chiodini
Travel Health Specialist Nurse

238

New immunisation training guidance



Revised and updated version of the original standards published by the former HPA in 2005. Previously published as two separate documents, this revised document incorporates the Minimum Standards and the Core Curriculum in one document.

The joint RCN/PHE documents 'Supporting the delivery of immunisation education', and the 'Immunisation knowledge and skills competence assessment tool' previously published by the RCN have both now been incorporated into this version.

Practitioners may also require additional training depending on the vaccine(s) they give. **For example, those who give travel immunisations will require specific training on travel health – a generic immunisation course alone would not be sufficient.**

Jane Chiodini
Travel Health Specialist Nurse

239

At the current time, Nursing Associates and Physician's Associates can only work under a PSD (the legislation does not allow them to work under a PGD).

There is no mention in the document in relation to travel vaccines but there is reference to National Minimum Standards

Practitioners may also require additional training depending on the vaccine(s) they give. For example, those who give travel immunisations will require specific training on travel health – a generic immunisation course alone would not be sufficient.

<https://www.rcn.org.uk/professional-development/qualifications/2017-2018>
<https://www.nmfs.org.uk/documents/ndocuments/ndocuments/standards/nursing-associates-proficiency-standards.pdf>
<https://www.nmfs.org.uk/documents/ndocuments/ndocuments/standards/physician-associates-proficiency-standards.pdf>

Jane Chiodini
 Trustee and Director

240

Intended to be free of charge for all

Access our e-learning programmes on the e-LfH Hub [Register / Log in](#)

e-LfH
 e-Learning for Health

Health Education England

Home Programmes About News Support Demo Contact us Search this website

Sections

Immunisation

An interactive e-learning programme to support the training of healthcare professionals involved in advising on and/or delivering immunisations across the life course.

This programme is in partnership with...

<https://www.e-lfh.org.uk/programmes/immunisation/>

Jane Chiodini
 Trustee and Director

241

Yellow card

YellowCard
 Reporting to MHRA

A side effect to a medicine?
 You can report it using **YellowCard**

Visit www.mhra.gov.uk/yellowcard to report suspected side effects

You can get Yellow Card forms:
 • from pharmacies or GP surgeries
 • by calling 0800 100 3352

Yellow Card - MHRA

Download

<https://yellowcard.mhra.gov.uk/downloadable-information/>

Jane Chiodini
 Trustee and Director

242

Licensed, Unlicensed and Off-label

NHS

Why is my child being given an 'off-label' vaccine?

A guide for parents*

Off-label vaccines

An introductory guide for healthcare professionals

The use of vaccines that have been temporarily stored outside the recommended temperature range

A brief guide for parents

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/390117/PHE_9373_VU_223_Dec_2014_11.pdf

Jane Chiodini
 Trustee and Director

243

Immunisation training includes...

- CPR and anaphylaxis
- Consent
- Prescribing
- Administration
- Documentation
- Vaccine storage – protocol
- Finance

Not part of the National Standards, but topics covered briefly on day 2

Jane Chiodini
 Trustee and Director

244

www.resus.org.uk

Resuscitation Council (UK)

HOME Guidelines Courses Statements Clinical info Publications A to Z index

search

LIFESAVER
 A free crisis simulator to teach resuscitation skills

Current Guidelines Quality Standards Donate

Annual training for CPR and anaphylaxis should be undertaken

Jane Chiodini
 Trustee and Director

245

Vaccine ordering, storage and handling




The perfect fridge!

Jane Chiodini
Travel Health Specialist Nurse

246

Prescribing for travel medicine

Legal requirement – covered on day 2




https://www.nice.org.uk/guidance/mpg2

Jane Chiodini
Travel Health Specialist Nurse

247

FAQs (under News) for information on prescribing




<http://www.janechiodini.co.uk/news/faqs/faq-no-1/>

Jane Chiodini
Travel Health Specialist Nurse

248

Consent




Chapter 2 in the 'Green Book'

Jane Chiodini
Travel Health Specialist Nurse

249

Administration

- Cleansing the skin
- Size of needles
- Preparing the vaccine
- Post vaccination waiting time ?



Jane Chiodini
Travel Health Specialist Nurse

250

Equipment



Jane Chiodini
Travel Health Specialist Nurse

251

Importance of Documentation – working within your code

The screenshot shows the NMC website with a focus on 'Hearings and outcomes'. A blue book titled 'The Code' is shown next to the website content. The website text includes: 'The schedule and results of hearings and the sanctions we've imposed' and 'Restrictions and sanctions: The different types of sanctions we can impose on a nurse or midwife'.

<http://www.nmc-uk.org/>

Jane Chiodini
Travel Health Specialist Nurse

252

Medicine Management

Standards for medicines management

These standards were withdrawn on 29 January 2019

The Standards for medicines management (2007) and underpinning NMC Circulars 08/2008 and 09/2008 were withdrawn on 29 January 2019. We did this because it's not within our remit as a regulator to provide the type of clinical practice guidance. However, we recognise that it's important that all healthcare professionals can access accurate information on the safe and effective handling, management and administration of medicines. This page has guidance on where you may be able to find this information.

Where to find information on medicines management

- Professional guidance on the safe and correct handling of medicines - we've worked closely with the Royal Pharmaceutical Society (RPS) and their stakeholder partners on this guidance for all healthcare professionals covering areas such as the storage, transportation and disposal of medicines.
- Professional guidance on the administration of medicines in healthcare settings - this guidance, co-produced by the Royal Pharmaceutical Society (RPS) and Royal College of Nursing (RCN), provides principles-based guidance to ensure the safe administration of medicines by healthcare professionals.
- Advisory guidance on administration of medicines by nursing associates - Health Education England (HEE) guidance.

Jane Chiodini
Travel Health Specialist Nurse

253

Resources for a travel service

The collage includes various resources such as 'Teaching tools', 'Key reference for best practice', 'Some key tools to consider the best', 'Key forms for risk assessment and analysis for your travellers', and 'Some sites to have handy'. A prominent poster for 'PLATFORM 9 3/4' is also visible.

Jane Chiodini
Travel Health Specialist Nurse

254

Agenda

- Introduction to travel medicine
- Travel risk assessment
- Travel vaccines and related issues
- Travel medicine operational issues
- Recap on resources

Jane Chiodini
Travel Health Specialist Nurse

255

Resources in Travel Health

The illustration shows three white, stylized human figures standing together, supporting a large, glowing globe of the Earth on their shoulders.

Jane Chiodini
Travel Health Specialist Nurse

256

Resources from my website www.janechiodini.co.uk

Jane Chiodini
Travel Health Specialist Nurse

Welcome

When I originally established my website I wanted to provide healthcare professionals with essential Travel Medicine information and resources. Thanks to improvements in technology and the range of resources available I'm happy to say there is now a wealth of accessible information. The aim now is to provide a "one stop shop" for information for travel health consultations. However I didn't realise what travel medicine is all about, click on the image below for a learning on the "Back Link" section at the bottom of this homepage. This will take you to a short module entitled "What is Travel". This will explain more about the field of practice!

Jane Chiodini
Travel Health Specialist Nurse

257

UK National databases - www.travax.nhs.uk and for the public - www.fitfortravel.nhs.uk

Jane Chiodini
Travel Health Specialist Nurse

258

<https://travelhealthpro.org.uk/>

From NaTHNaC for healthcare professionals and the general public

Jane Chiodini
Travel Health Specialist Nurse

259

Key UK guidance resources

Both available from the publications page on my website
<http://www.janechiodini.co.uk/about/publications/>

Jane Chiodini
Travel Health Specialist Nurse

260

Jane Chiodini's Blog

Jane Chiodini
Travel Health Specialist Nurse

261

Jane Chiodini
Travel Health Specialist Nurse

262

Access via the 'New to Travel' page on my website


Please note, if using TRAVAX, you will still need a user name and password

<http://www.janechiodini.co.uk/tools/new-to-travel/>

Jane Chiodini
Travel Health Specialist Nurse

263

Key UK resources




- Department of Health 'Green Book' published in 2006 but new cover and new publication date of 2013
- DO NOT USE PAPER COPY
- Use online version
 - Whole book
 - Individual chapters
 - Update patches

Jane Chiodini
Thames Valley University

264

Key UK resources




- UK Malaria Guidelines found at www.malaria-reference.co.uk
- Also available from the malaria page on my website <http://www.janechiodini.co.uk/news/help/malaria/>

Jane Chiodini
Thames Valley University

265

DH, PHE and MHRA all under one roof




<https://www.gov.uk/government/organisations/department-of-health>
<https://www.gov.uk/government/organisations/public-health-england>

Jane Chiodini
Thames Valley University

266

Working within National Guidelines and knowing one's limitations



Telephone helplines

NaTHNaC
0845 602 6712
Monday – Friday 9am to 11.00 am and 1pm to 2pm x 2
Closed Monday and Friday at 2pm and other days at 3.30pm

TRAVAX
0141 300 1130
Mon. & Wed. 2 to 4pm
Friday 9.30 to 11.30am


MRL e mail service
Download risk assessment form from www.malaria-reference.co.uk, complete and return by e mail

Jane Chiodini
Thames Valley University

267

Electronic Medicines Compendium

www.medicines.org.uk – provides SmPCs and PILs



Don't forget the protected login area to store your own choices

Jane Chiodini
Thames Valley University

268

Immunisation training

National Minimum Standards and Core Curriculum for Immunisation Training for Registered Healthcare Practitioners

Revised February 2018



<https://www.janechiodini.co.uk/help/immunisation-resources/>

Jane Chiodini
Thames Valley University

269

Finding additional destinations


Google search www.google.co.uk
and/or google maps
<http://maps.google.co.uk/>



Jane Chiodini
Travel Health Specialist Nurse

270

Worldwide resources




TRAVAX from Shoreland is not the same as UK TRAVAX

Jane Chiodini
Travel Health Specialist Nurse

271

Other useful maps also at www.who.int/ith



http://gamapservr.who.int/maplibrary/Files/Maps/Global_CholeraCases_IJTRiskMap.png

Jane Chiodini
Travel Health Specialist Nurse

272

World Health Organization www.who.int

or Google title of the WHO Factsheet required

an extremely useful website with so much information e.g.

Substandard and falsified medical products



Jane Chiodini
Travel Health Specialist Nurse

273

Finding travel clinics abroad www.istm.org



Jane Chiodini
Travel Health Specialist Nurse

274

Additional websites and resources on my website



Jane Chiodini
Travel Health Specialist Nurse

275

**Interim work between the two study days
listed on your webpage**

- Play the vaccine videos to recap on this aspect
- Look around my website
- Find out what travel PGDs you have at work
- See if you have a vaccine storage protocol at work
- Do the practice case study e learning on your page if you have time left
- Remember you can go into the e learning to reflect on today's presentation if you wish – but only available for a limited time.

Jane Chiodini
Theobald Group

276

**Don't forget to view the tasks suggested if you
can for the interim period.
See you on Monday 30th September 2019**



Jane Chiodini
Theobald Group

277